

DIATER Parenteral

Bacterial suspension of germ culture. For specific immunotherapy.

COMPOSITION

Sterile suspension of inactivated germs in a 0.3% phenolated normal saline solution, at the medically prescribed concentration, originating from the culture of the sample the physician took from the patient that is sent to the laboratory for processing, or that which the specialist decides.

INDICATIONS

Rhinitis, sinusitis, otitis, pharyngitis, laryngitis, bronchitis, conjunctivitis or other delayed hypersensitivity ailments.

HOW SUPPLIED

Vial no.	Label Colour	Volume
0	Grey	1.5 mL
1	Green	4.5 mL
2	Yellow	4.5 mL
3	Red	5 mL

INITIAL TREATMENT

V3: Container with four vials (0-1-2-3)

MAINTENANCE TREATMENT

V4: Container with two vials 3

Initial treatment:

Vial (colour/no.)	Injection Volume	Dosage Frequency
0	0.5 mL	4 days
1	0.1 mL	4 days
	0.2 mL	4 days
	0.4 mL	4 days
	0.6 mL	4 days
	0.8 mL	4 days
	1.0 mL	4 days
2	0.1 mL	4 days
	0.2 mL	4 days
	0.4 mL	4 days
	0.6 mL	4 days
	0.8 mL	4 days
	1.0 mL	4 days
3	0.1 mL	4 days
	0.2 mL	4 days
	0.4 mL	4 days
	0.6 mL	4 days
	0.8 mL	4 days
	1.0 mL	4 days

Maintenance treatment:

Vial (colour/no.)	Injection Volume	Dosage Frequency
3	1 mL	2 weeks
	1 mL	2 weeks
	1 mL	2 weeks
	1 mL	2 weeks
3	1 mL	2 weeks
	1 mL	2 weeks
	1 mL	2 weeks
	1 mL	2 weeks

ROUTE OF ADMINISTRATION

Subcutaneous injection

RECOMMENDED ADMINISTRATION PLAN

The plan recommended by the laboratory is subject to whatever modification the specialist deems necessary.