

DIATER Nasopharyngeal

Specific bacterial suspension of germ culture for spray. For specific immunotherapy.

COMPOSITION

Sterile suspension of inactivated germs in a 0.3% phenolated normal saline solution, in a concentration of 500 MG/mL (million germs per mL) vial **A**, and 1.000 MG/mL (million germs per mL) vial **B**, or as prescribed by the physician, originating from the culture of the sample the physician took from the patient that is sent to the laboratory for processing, or that which the specialist decides.

INDICATIONS

Rhinitis, sinusitis, otitis, laryngitis, bronchitis, conjunctivitis, bronchial asthma, or other delayed hypersensitivity ailments.

HOW SUPPLIED

Vial letter	Label Colour	Volume
A	Yellow	10 mL
B	Red	10 mL

INITIAL TREATMENT

V1: Container with two vials (A and B), 10 mL/vial.

MAINTENANCE TREATMENT

V8: Container with 1 vial (B), 10 mL/vial.

V6: Container with 2 vials (B), 10 mL/vial.

Initial treatment:

Vial (colour/letter)	Dose	Dosage Frequency
A	1 Actuation in each nostril and pharynx until finishing the contents of the vial.	Daily
B	1 Actuation in each nostril and pharynx until finishing the contents of the vial.	Daily

Maintenance treatment:

Vial (colour/letter)	Dose	Dosage Frequency
B	1 Actuation in each nostril and pharynx until finishing the contents of the vial.	Daily
B	1 Actuation in each nostril and pharynx until finishing the contents of the vial.	Daily

ROUTE OF ADMINISTRATION

Nasopharyngeal.

RECOMMENDED DOSAGE PLAN

The plan recommended by the laboratory is subject to whatever modification the specialist deems necessary.